

# COSE215: Theory of Computation

## Lecture 8 — Properties of Regular Languages (2):

### **Pumping Lemma**

Hakjoo Oh  
2018 Spring

## Some Fundamental Questions

So far, we have studied regular languages. But, some fundamental questions remain:

- Are all languages regular?
  - ▶ No, e.g.,  $L = \{a^n b^n \mid n \geq 0\}$  is not regular.
- How to prove that a language is non-regular? Two methods:
  - 1 Direct proof by Pigeonhole principle.
  - 2 By using the pumping lemma.

## Example 1: $L = \{a^n b^n \mid n \geq 0\}$ is non-regular

Direct proof:

- Proof by contradiction.
- The basic tool: Pigeonhole principle: If you put more than  $n$  pigeons into  $n$  holes, then some hole has more than one pigeon.
- Assume  $L$  is regular.
- Then there is a DFA  $M = (Q, \Sigma, \delta, q_0, F)$  recognizing  $L$ .
- Define:
  - ▶ Pigeons =  $\{a^n \mid n \geq 0\} = \{a, aa, aaa, \dots\}$
  - ▶ Holes = states in  $Q$
- Put pigeon  $a^n$  into hole  $\delta^*(q_0, a^n)$ 
  - ▶ i.e., the hole corresponding to the state reached by input  $a^n$
- We have  $|Q|$  holes but more than  $|Q|$  pigeons (actually, infinitely many).
- So, two pigeons must be put in the same hole, say  $a^i$  and  $a^j$ , where  $i \neq j$ .
  - ▶ That is,  $a^i$  and  $a^j$  lead to the same state.
- Then, since  $M$  accepts  $a^i b^i$ , it also accepts  $a^j b^i$ , which is a contradiction.
- Thus, the original assumption that  $L$  is regular is false,
- That is,  $L$  is non-regular.

## Example 2: $L = \{ww \mid w \in \{0, 1\}^*\}$ is non-regular

- Show by contradiction, using Pigeonhole principle.
- Assume  $L$  is regular, so there is a DFA  $M = (Q, \Sigma, \delta, q_0, F)$  recognizing  $L$ .
- Define:
  - ▶ Pigeons =  $\{0^i1 \mid i \geq 0\} = \{1, 01, 001, \dots\}$
  - ▶ Holes = states in  $Q$
- Put pigeon string  $0^i1$  into hole  $\delta^*(q_0, 0^i1)$
- By Pigeonhole principle, two pigeons share a hole, say  $0^i1$  and  $0^j1$ , where  $i \neq j$ .
- So  $0^i1$  and  $0^j1$  lead to the same state.
- $M$  accepts  $0^i10^i1$ , so does  $0^j10^i1$ , which is a contradiction.

# The Pumping Lemma

## Theorem (Pumping Lemma)

For any regular language  $L$  *there exists* an integer  $n$ , such that *for all*  $x \in L$  with  $|x| \geq n$ , *there exist*  $u, v, w \in \Sigma^*$ , such that

- 1  $x = uvw$
- 2  $|uv| \leq n$
- 3  $|v| \geq 1$
- 4 *for all*  $i \geq 0$ ,  $uv^i w \in L$ .

# Proof of the Pumping Lemma

- Let  $M$  be a DFA for  $L$ . Suppose  $M$  has  $n$  states.
- Take  $x \in L$  with  $|x| \geq n$ , let  $m = |x|$ :

$$x = a_1 a_2 \dots a_m$$

- Let  $p_i = \delta^*(q_0, a_1 a_2 \dots a_i)$ . Note  $p_0 = q_0$  and  $p_m$  is a final state.
- Consider the first  $n + 1$  states:  $p_0 p_1 \dots p_n$ .
- By Pigeonhole principle, two  $p_i$  and  $p_j$  with  $0 \leq i < j \leq n$  share a state, i.e.,  $p_i = p_j$ .
- Break  $x = uvw$ :
  - ▶  $u = a_1 a_2 \dots a_i$
  - ▶  $v = a_{i+1} a_{i+2} \dots a_j$
  - ▶  $w = a_{j+1} a_{j+2} \dots a_m$
- Note that  $\delta^*(p_0, u) = p_i$ ,  $\delta^*(p_i, v) = p_i$ , and  $\delta^*(p_i, w) = p_m$ .
- Thus,  $\delta^*(p_0, uv) = p_m$ ,  $\delta^*(p_0, uvw) = p_m$ ,  $\delta(p_0, uv^2w) = p_m$ , and so on.

## Using Pumping Lemma to show non-regularity

- If  $L$  is regular,  $L$  satisfies pumping lemma?
- If  $L$  satisfies pumping lemma,  $L$  is regular?
- If  $L$  does not satisfy pumping lemma, then  $L$  is non-regular?

Pumping lemma can be used only for proving languages not to be regular.

## Example 1

Prove that  $L = \{0^i 1^i \mid i \geq 0\}$  is not regular.

- Show that pumping lemma (P.L.) does not hold.
- If  $L$  is regular, then by P.L. there exists  $n$  such that ...
- Now let  $x = 0^n 1^n$
- $x \in L$  and  $|x| \geq n$ , so by P.L. there exist  $u, v, w$  such that (1)–(4) hold.
- We show that for all  $u, v, w$  (1)–(4) do not all hold.
- If (1), (2), (3) hold then  $x = 0^n 1^n = uvw$  with  $|uv| \leq n$  and  $|v| \geq 1$ .
- So,  $u = 0^s, v = 0^t, w = 0^p 1^n$  with

$$s + t \leq n, \quad t \geq 1, \quad p \geq 0, \quad s + t + p = n.$$

- Then (4) fails for  $i = 0$ :

$$uv^0w = uw = 0^s 0^p 1^n = 0^{s+p} 1^n \notin L, \quad \text{since } s + p \neq n$$



## Example 2

Prove that  $L = \{ww^R \mid w \in \{a, b\}^*\}$  is not regular.

- Show that pumping lemma (P.L.) does not hold.
- If  $L$  is regular, then by P.L. there exists  $n$  such that ...
- Now let  $x = a^n b^n b^n a^n$
- $x \in L$  and  $|x| \geq n$ , so by P.L. there exist  $u, v, w$  such that (1)–(4) hold.
- We show that for all  $u, v, w$  (1)–(4) do not all hold.
- If (1), (2), (3) hold then  $x = a^n b^n b^n a^n = uvw$  with  $|uv| \leq n$  and  $|v| \geq 1$ .
- So,  $u = a^s, v = a^t, w = a^p b^n b^n a^n$  with

$$s + t \leq n, \quad t \geq 1, \quad p \geq 0, \quad s + t + p = n.$$

- Then (4) fails for  $i = 0$ :

$$uv^0w = uw = a^s a^p b^n b^n a^n = a^{s+p} b^n b^n a^n \notin L, \text{ since } s + p \neq n$$

## Example 3

Prove that  $L = \{w \in \{a, b\}^* \mid n_a(w) < n_b(w)\}$  is not regular.

- Show that pumping lemma (P.L.) does not hold.
- If  $L$  is regular, then by P.L. there exists  $n$  such that ...
- Now let  $x = a^n b^{n+1}$
- $x \in L$  and  $|x| \geq n$ , so by P.L. there exist  $u, v, w$  such that (1)–(4) hold.
- We show that for all  $u, v, w$  (1)–(4) do not all hold.
- If (1), (2), (3) hold then  $x = a^n b^{n+1} = uvw$  with  $|uv| \leq n$  and  $|v| \geq 1$ .
- So,  $u = a^s, v = a^t, w = a^p b^{n+1}$  with

$$s + t \leq n, \quad t \geq 1, \quad p \geq 0, \quad s + t + p = n.$$

- Then (4) fails for  $i = 2$ :

$$uv^2w = a^s a^{2t} a^p b^{n+1} = a^{s+2t+p} b^{n+1} \notin L,$$

since  $s + 2t + p \geq n + 1$ .

## Example 4

Prove that  $L = \{a^n \mid n \text{ is a perfect square}\}$  is not regular.

- Show that pumping lemma (P.L.) does not hold.
- If  $L$  is regular, then by P.L. there exists  $n$  such that ...
- Now let  $x = a^{n^2}$
- $x \in L$  and  $|x| \geq n$ , so by P.L. there exist  $u, v, w$  such that (1)–(4) hold.
- We show that for all  $u, v, w$  (1)–(4) do not all hold.
- If (1), (2), (3) hold then  $x = a^{n^2} = uvw$  with  $|uv| \leq n$  and  $|v| \geq 1$ .
- Then, clearly  $v = a^k$  with  $1 \leq k \leq n$ .
- Then (4) fails for  $i = 0$ :

$$uv^0w = a^{n^2-k} \notin L, \quad \text{since } n^2 - k > (n - 1)^2.$$

## Example 5

Prove that  $L = \{a^n b^k c^{n+k} \mid n \geq 0 \wedge k \geq 0\}$  is not regular.

- It is not difficult to apply the pumping lemma directly, but it is even easier to use closure under homomorphism. Take

$$h(a) = a, \quad h(b) = a, \quad h(c) = c,$$

then

$$h(L) = \{a^{n+k} c^{n+k} \mid n + k \geq 0\} = \{a^i b^i \mid i \geq 0\}.$$

We know this language is not regular.

- Also, we know that if a language  $L_1$  is regular, then  $h(L_1)$  is regular. Taking its contraposition, we conclude that  $L$  is not regular.

## cf) The converse of pumping lemma is not true

$$L = \{c^m a^n b^n \mid m \geq 1, n \geq 1\}$$

- $L$  satisfies the pumping lemma.
  - ▶ For any  $x \in L$  of length  $\geq 1$ , we can take  $u = \epsilon$ ,  $v =$  the first letter of  $x$  ( $c$ ), and  $w =$  the rest of  $x$ .
- However,  $L$  is not regular.
  - ▶ We can prove this using a general version of pumping lemma: For any regular language  $L$ , there exists  $n \geq 1$  such that for every string  $uvw \in L$  with  $|w| \geq p$  such that
    - ★  $uvw = uxyzv$
    - ★  $|xy| \leq n$
    - ★  $|y| \geq 1$
    - ★ For all  $i \geq 0$ ,  $uxy^i z v \in L$ .
- Still, the converse of the general lemma is not true.
  - ▶ Languages that satisfy the lemma can still be non-regular.
  - ▶ For a necessary and sufficient condition to be regular, refer to Myhill-Nerode theorem.