

COSE212: Programming Languages

Lecture 1 — Inductive Definitions (1)

Hakjoo Oh
2018 Fall

Inductive Definitions

Inductive definition (induction) is widely used in the study of programming languages and computer science in general: e.g.,

- The syntax and semantics of programming languages
- Data structures (e.g., lists, trees, graphs)

Induction is a technique for formally defining a set:

- The set is defined in terms of itself.
- The only way of defining an infinite set by a finite means.

Three styles to inductive definition:

- Top-down
- Bottom-up
- Rules of inference

Example (Top-Down)

Let us define a certain subset S of natural numbers (\mathbb{N}) as follows:

Definition (S)

A natural number n is in S if and only if

- 1 $n = 0$, or
- 2 $n - 3 \in S$.

The definition is *inductive*, because the set is defined in terms of itself.
What is the set S ?

Example (Continued)

Let us see what natural numbers are in S .

- 0 is in S because of the first condition of the definition.
- 3 is in S because $3 - 3 = 0$ and 0 is in S .
- 6 is in S because $6 - 3 = 3$ and 3 is in S .
- ...

We can conjecture that $\{0, 3, 6, 9, \dots\} \subseteq S$.

Proof by mathematical induction .

We show that $3k \in S$ for all $k \in \mathbb{N}$.

- 1 Base case: $3k \in S$ when $k = 0$.
- 2 Inductive case: Assume $3k \in S$ (Induction Hypothesis, I.H.).
Then show $3 \cdot (k + 1) \in S$, which holds because
 $3 \cdot (k + 1) - 3 = 3k \in S$ by the induction hypothesis.



Example (Continued)

What about other numbers? Does S contain only the multiples of 3 ?

- For instance, $1 \in S$? No. Because the first condition is not true, the second condition must be true for 1 to be in S . However, it is not true because $1 - 3 = -2$ is not a natural number. Similarly, we can show that $2 \notin S$.
- What about 4 ? Because $4 - 3 = 1 \notin S$, $4 \notin S$.

By similar reasoning, we can conjecture that if n is not a multiple of 3 then n is not in S . In other words, S contains multiples of 3 only: i.e.,

$$\{0, 3, 6, 9, \dots\} \supseteq S.$$

Proof by contradiction.

Let $n = 3k + q$ ($q = 1$ or 2) and assume $n \in S$. By the definition of S , $n - 3, n - 6, \dots, n - 3k \in S$. Thus, S must include 1 or 2 , a contradiction. □

A Bottom-up Definition

An alternative inductive definition of S :

Definition (S)

S is the *smallest* set such that $S \subseteq \mathbb{N}$ and S satisfies the following two conditions:

- 1 $0 \in S$, and
- 2 if $n \in S$, then $n + 3 \in S$.

- The two conditions imply $\{0, 3, 6, 9, \dots\} \subseteq S$.
- The two conditions do not imply $\{0, 3, 6, 9, \dots\} \supseteq S$. E.g.,
 - ▶ \mathbb{N} satisfies the conditions: $0 \in \mathbb{N}$ and if $n \in \mathbb{N}$ then $n + 3 \in \mathbb{N}$.
 - ▶ $\{0, 3, 6, 9, \dots\} \cup \{1, 4, 7, 10, \dots\}$ satisfies the conditions.
- This is why the definition requires S to be the **smallest** such a set.
- The smallest set that satisfies the two conditions is unique:

$$S = \{0, 3, 6, 9, \dots\}.$$

Rules of Inference

The third way is to define the set with inference rules. An inference rule is of the form:

$$\frac{A}{B}$$

- A : hypothesis (antecedent)
- B : conclusion (consequent)
- “if A is true then B is also true”.
- \overline{B} : axiom (inference rule without hypothesis)

The hypothesis may contain multiple statements:

$$\frac{A \quad B}{C}$$

“If both A and B are true then so is C ”.

Rules of Inferences

The set \mathcal{S} is defined as inference rules as follows:

Definition (\mathcal{S})

$$\overline{0 \in \mathcal{S}} \quad \frac{n \in \mathcal{S}}{(n + 3) \in \mathcal{S}}$$

Interpret the rules as follows:

“A natural number n is in \mathcal{S} iff $n \in \mathcal{S}$ can be derived from the axiom by applying the inference rules finitely many times”

For example, $3 \in \mathcal{S}$ because we can find a “proof/derivation tree”:

$$\overline{0 \in \mathcal{S}} \text{ the axiom}$$
$$\overline{3 \in \mathcal{S}} \text{ the second rule}$$

but $1, 2, 4, \dots \notin \mathcal{S}$ because we cannot find proofs. Note that this interpretation enforces that \mathcal{S} is the smallest set closed under the inference rules.

Exercises

- ① What set is defined by the following inductive rules?

$$\overline{\mathbf{3}} \quad \frac{x \quad y}{x + y}$$

- ② What set is defined by the following inductive rules?

$$\overline{()} \quad \frac{s}{(s)} \quad \frac{s}{s \ s}$$

- ③ Define the following set as rules of inference:

$$S = \{a, b, aa, ab, ba, bb, aaa, aab, aba, abb, baa, bab, bba, bbb, \dots\}$$

- ④ Define the following set as rules of inference:

$$S = \{x^n y^{n+1} \mid n \in \mathbb{N}\}$$

Summary

In inductive definitions, a set is defined in terms of itself. Three styles:

- Top-down
- Bottom-up
- Rules of inference

In PL, we mainly use the rules-of-inference method.