

COSE215: Theory of Computation

Lecture 3 — Nondeterministic Finite Automata

Hakjoo Oh
2018 Spring

Definition

Definition (NFA)

A *nondeterministic finite automaton* (or *NFA*) is defined as,

$$M = (Q, \Sigma, \delta, q_0, F)$$

where

- Q : a finite set of *states*
- Σ : a finite set of *input symbols* (or input alphabet)
- $q_0 \in Q$: the *initial state*
- $F \subseteq Q$: a set of *final states*
- $\delta : Q \times \Sigma \rightarrow 2^Q$: *transition function*


Example

$$(\{q_0, q_1, q_2\}, \{0, 1\}, \delta, q_0, \{q_2\})$$


$$\delta(q_0, 0) = \{q_0, q_1\} \quad \delta(q_0, 1) = \{q_0\}$$

$$\delta(q_1, 0) = \emptyset \quad \delta(q_1, 1) = \{q_2\}$$

$$\delta(q_2, 0) = \emptyset \quad \delta(q_2, 1) = \emptyset$$



cf) Compare with the equivalent DFA:




Extended Transition Function

$$\delta^* : Q \times \Sigma^* \rightarrow 2^Q$$

- (Basis) $s = \epsilon$:
- (Induction) $s = wa$:

Example



$$\delta^*(q_0, 00101) = \bigcup_{s_i \in \delta^*(q_0, 0010)} \delta(s_i, 1) = \delta(q_0, 1) \cup \delta(q_1, 1) = \{q_0\} \cup \{q_2\} = \{q_0, q_2\}$$

$$\delta^*(q_0, 0010) = \bigcup_{s_i \in \delta^*(q_0, 001)} \delta(s_i, 0) = \delta(q_0, 0) \cup \delta(q_2, 0) = \{q_0, q_1\} \cup \emptyset = \{q_0, q_1\}$$

$$\delta^*(q_0, 001) = \bigcup_{s_i \in \delta^*(q_0, 00)} \delta(s_i, 1) = \delta(q_0, 1) \cup \delta(q_1, 1) = \{q_0\} \cup \{q_2\} = \{q_0, q_2\}$$

$$\delta^*(q_0, 00) = \bigcup_{s_i \in \delta^*(q_0, 0)} \delta(s_i, 0) = \delta(q_0, 0) \cup \delta(q_1, 0) = \{q_0, q_1\} \cup \emptyset = \{q_0, q_1\}$$

$$\delta^*(q_0, 0) = \bigcup_{s_i \in \delta^*(q_0, \epsilon)} \delta(s_i, 0) = \delta(q_0, 0) = \{q_0, q_1\}$$

$$\delta^*(q_0, \epsilon) = \{q_0\}$$

Exercise: Language of an NFA

The language of NFA $M = (Q, \Sigma, \delta, q_0, F)$ is defined as follows:

$$L(M) = \{ \quad \quad \quad \}$$

Exercises

Design NFAs for the following languages:

- 1 $L = \{a^n b \mid n \geq 0\}$
- 2 $L = \{x01y \mid x, y \in \{0, 1\}^*\}$
- 3 $L = \{01w \mid w \in \{0, 1\}^*\}$
- 4 $L = \{w \in \{0, 1\}^* \mid w \text{ contains at least two } 0\text{'s}\}$
- 5 $L = \{w \in \{0, 1\}^* \mid w \text{ contains exactly two } 0\text{'s}\}$
- 6 $L = \{w \in \{0, 1\}^* \mid w \text{ has three consecutive } 0\text{'s}\}$

Equivalence of DFA and NFA

Theorem (Equivalence)

A Language L is accepted by some NFA if and only if L is accepted by some DFA.

Proof.

By the two Lemmas below. □

Lemma (DFA to NFA)

Given a DFA D , there always exists an NFA N such that $L(D) = L(N)$.

Lemma (NFA to DFA)

Given an NFA N , there always exists a DFA D such that $L(N) = L(D)$.

DFA to NFA

Lemma (DFA to NFA)

Given a DFA D , there always exists an NFA N such that $L(D) = L(N)$.

Proof) Assume a DFA $D = (Q, \Sigma, \delta_D, q_0, F)$ is given. Define an NFA as follows:

$$N = (Q, \Sigma, \delta_N, q_0, F) \text{ where } \delta_N(q, a) = \{\delta_D(q, a)\}$$

To prove:

$$L(D) = \{w \in \Sigma^* \mid \delta_D^*(q_0, w) \in F\} = \{w \in \Sigma^* \mid \delta_N^*(q_0, w) \cap F \neq \emptyset\} = L(N)$$

It is enough to show that

$$\delta_N^*(q_0, w) = \{\delta_D^*(q_0, w)\}$$

The proof is by induction on $|w|$.

- $w = \epsilon$: By the definitions of δ_D^* and δ_N^* , $\delta_D^*(q_0, \epsilon) = q_0$ and $\delta_N^*(q_0, \epsilon) = \{q_0\}$.
- $w = sa$:

$$\begin{aligned} \delta_N^*(q_0, sa) &= \bigcup_{s_i \in \delta_N^*(q_0, s)} \delta_N(s_i, a) && \text{by definition of } \delta_N^* \\ &= \delta_N(\delta_D^*(q_0, s), a) && \text{by I.H.} \\ &= \{\delta_D(\delta_D^*(q_0, s), a)\} && \text{by definition of } \delta_N \\ &= \{\delta_D^*(q_0, sa)\} && \text{by definition of } \delta_D^* \end{aligned}$$

NFA to DFA (Subset Construction)

Lemma (NFA to DFA)

Given an NFA N , there always exists a DFA D such that $L(N) = L(D)$.

Proof) Assume an NFA $N = (Q_N, \Sigma, \delta_N, q_0, F_N)$. Define a DFA as follows

$$D = (Q_D, \Sigma, \delta_D, \{q_0\}, F_D)$$

where

- $Q_D = 2^{Q_N}$
- $F_D = \{S \in Q_D \mid S \cap F_N \neq \emptyset\}$.
- For each $S \in Q_D$ and input symbol $a \in \Sigma$:

$$\delta_D(S, a) = \bigcup_{p \in S} \delta_N(p, a)$$

NFA to DFA

Then, we can prove $L(N) = L(D)$ by showing that

$$\delta_D^*({q_0}, w) = \delta_N^*(q_0, w).$$

The proof is by induction on the length of w .

- $w = \epsilon$: By definition, $\delta_D^*({q_0}, \epsilon) = \{q_0\} = \delta_N^*(q_0, \epsilon)$.
- $w = sa$: Induction hypothesis (I.H.):

$$\delta_D^*({q_0}, s) = \delta_N^*(q_0, s).$$

$$\begin{aligned} \delta_D^*({q_0}, sa) &= \delta_D(\delta_D^*({q_0}, s), a) && \text{by definition of } \delta_D^* \\ &= \delta_D(\delta_N^*(q_0, s), a) && \text{by I.H.} \\ &= \bigcup_{p \in \delta_N^*(q_0, s)} \delta_N(p, a) && \text{by definition of } \delta_D \\ &= \delta_N^*(q_0, sa) && \text{by definition of } \delta_N^* \end{aligned}$$



Example: Subset Construction


Find a DFA that is equivalent to:

$$N = (\{q_0, q_1, q_2\}, \{0, 1\}, \delta, q_0, \{q_2\})$$

$$\delta(q_0, 0) = \{q_0, q_1\} \quad \delta(q_0, 1) = \{q_0\}$$

$$\delta(q_1, 0) = \emptyset \quad \delta(q_1, 1) = \{q_2\}$$

$$\delta(q_2, 0) = \emptyset \quad \delta(q_2, 1) = \emptyset$$



Example: Subset Construction

$$D = (Q_D, \{0, 1\}, \delta_d, \{q_0\}, F_D)$$

- $Q_D = 2^{\{q_0, q_1, q_2\}} = \{\emptyset, \{q_0\}, \{q_1\}, \dots, \{q_0, q_1, q_2\}\}$
- $F_D = \{\{q_2\}, \{q_0, q_2\}, \{q_1, q_2\}, \{q_0, q_1, q_2\}\}$
- δ_D :

	0	1
\emptyset	\emptyset	\emptyset
$\rightarrow \{q_0\}$	$\{q_0, q_1\}$	$\{q_0\}$
$\{q_1\}$	\emptyset	$\{q_2\}$
$*\{q_2\}$	\emptyset	\emptyset
$\{q_0, q_1\}$	$\{q_0, q_1\}$	$\{q_0, q_2\}$
$*\{q_0, q_2\}$	$\{q_0, q_1\}$	$\{q_0\}$
$*\{q_1, q_2\}$	\emptyset	$\{q_2\}$
$*\{q_0, q_1, q_2\}$	$\{q_0, q_1\}$	$\{q_0, q_2\}$

Example: Subset Construction

