

COSE215: Theory of Computation

Lecture 1 — Mathematical Preliminaries

Hakjoo Oh
2016 Spring

Today

- Icebreaking: Introduce yourself
- Mathematical backgrounds and notation
 - ▶ Sets
 - ▶ Inductive proofs

Icebreaking

Introduce yourself:

- Free format. Say anything.
- Nothing to talk about? major, grade, interests, hobbies, specialty, goal, motivation for this course, what you expect from this course, etc

Sets

- A set is a collection of elements, e.g.,
 - ▶ $S = \{0, 1, 2\} = \{x \in \mathbb{N} \mid 0 \leq x \leq 2\}$
 - ▶ $S = \{2, 4, 6, \dots\} = \{x \in \mathbb{N} \mid x \text{ is even}\}$
- Notations:
 - ▶ \emptyset : the empty set
 - ▶ $S_1 \subseteq S_2$ iff $\forall x \in S_1. x \in S_2$
 - ▶ $S_1 \subset S_2$ if $S_1 \subseteq S_2$ and $S_1 \neq S_2$, e.g., $\{1, 2\} \subset \{1, 2, 3\}$, $\{1, 2\} \not\subset \{1, 2\}$
 - ▶ $|S|$: the number of elements in set S
 - ▶ S_1 and S_2 are disjoint iff $S_1 \cap S_2 = \emptyset$.

Construction of Sets

- Union, intersection, and difference:

$$S_1 \cup S_2 = \{x \mid x \in S_1 \vee x \in S_2\}$$

$$S_1 \cap S_2 = \{x \mid x \in S_1 \wedge x \in S_2\}$$

$$S_1 - S_2 = \{x \mid x \in S_1 \wedge x \notin S_2\}$$

- $\bar{S} = \{x \mid x \in U \wedge x \notin S\}$
- Powerset: $2^S = \mathcal{P}(S) = \{x \mid x \subseteq S\}$
- Cartesian product:

$$S_1 \times S_2 = \{(x, y) \mid x \in S_1 \wedge y \in S_2\}$$

In general,

$$S_1 \times S_2 \times \cdots \times S_n = \{(x_1, x_2, \dots, x_n) \mid x_i \in S_i\}$$

Partition

When S_1, S_2, \dots, S_n are subsets of a given set S , S_1, S_2, \dots, S_n forms a partition of S iff:

- 1 S_1, S_2, \dots, S_n are mutually disjoint:

$$\forall i, j. i \neq j \implies S_i \cap S_j = \emptyset$$

- 2 S_1, S_2, \dots, S_n cover S :

$$\bigcup_{1 \leq i \leq n} S_i = S$$

- 3 none of S_i is empty: $\forall i. S_i \neq \emptyset$.

Inductive proofs

In CS, every set is inductively defined. E.g.,

Example (Inductive Definition of Trees)

A set of trees is defined as follows:

- 1 (Basis) A single node (called root) is a tree.
- 2 (Induction) If T_1, T_2, \dots, T_k are trees, then the following is also a tree:
 - 1 Begin with a new node N , which is the root of the tree.
 - 2 Add edges from N to the roots of each of the trees T_1, T_2, \dots, T_k .

Example (Inductive Definition of Arithmetic Expressions)

A set of arithmetic expressions is defined as follows:

- (Basis) Any number or letter (i.e., a variable) is an expression.
- (Induction) If E and F are expressions, then so are $E + F$, $E * F$, and (E) .

Inductive Proofs

Induction is used to prove properties about inductively defined sets. Let S be an inductively-defined set. Let $P(x)$ be a property of x . To show that, for all $x \in S.P(x)$, it suffices to show that:

- 1 (Base case): Show $P(x)$ for all basis elements $x \in S$.
- 2 (Inductive case): For each inductive rule using elements x_1, \dots, x_k of S to construct an element x , show that

if $P(x_1), \dots, P(x_k)$ then $P(x)$

$P(x_1), \dots, P(x_k)$: induction hypotheses.

Inductive Proofs: Example

Prove that every tree has one more node than it has edges.

Proof.

Formally, what we prove is $P(T)$ = “if T is a tree, and T has n nodes and e edges, then $n = e + 1$ ”.

- 1 Base case: The base case is when T is a single node. Then, $n = 1$ and $e = 0$, so the relationship $n = e + 1$ holds.
- 2 Inductive case: The inductive case is when T is built with root node N and k smaller trees T_1, T_2, \dots, T_k .
 - 1 **Induction hypothesis:** The statements $P(T_i)$ holds for $i = 1, 2, \dots, k$. That is T_i have n_i nodes and e_i edges; then $n_i = e_i + 1$.
 - 2 **To Show:** $P(T)$ holds: if T has n nodes and e edges, then $n = e + 1$. The nodes of T are node N and all the nodes of the T_i 's, i.e., $n = 1 + n_1 + \dots + n_k$. The edges of T are the k edges we added explicitly in the inductive definition step, plus the edges of the T_i 's. Hence, T has $e = k + e_1 + \dots + e_k$ edges.

$$\begin{aligned}n &= 1 + n_1 + \dots + n_k && \text{def. of } n \\&= 1 + (e_1 + 1) + \dots + (e_k + 1) && \text{induction hypothesis} \\&= 1 + k + e_1 + \dots + e_k \\&= 1 + e && \text{def. of } e\end{aligned}$$

Inductive Proofs: Example

Prove that every expression has an equal number of left and right parentheses.

Proof.

Formally, the formal statement $P(G)$ we need to prove is: "if G has l left parentheses and r right parentheses, then $l = r$."

- 1 **Base case:** The base case is when G is a number or a variable, in which cases $l = r = 0$.
- 2 **Inductive case:** There are three cases, where G is constructed recursively from smaller expressions:

▶ $G = E + F$:

- 1 **Induction hypothesis:** The statement holds for all smaller expressions: for E , $l_E = r_E$, and for F , $l_F = r_F$.
- 2 **To Show:** $P(G)$ holds: $l_G = r_G$:

$$\begin{aligned}l_G &= l_E + l_F \\ &= r_E + r_F \quad \text{I.H.} \\ &= r_G\end{aligned}$$

- ▶ $G = E * F$: similar
- ▶ $G = (E)$: similar



Summary

- Sets: definition, notations, constructions
- Inductive definitions and proofs.